

would our message need to change once more? Here are the facts: hepatitis C virus is common among Americans born between 1945 and 1965. A current Centers for Disease Control and Prevention factsheet asserts that most people infected with hepatitis C virus are unaware of their diagnosis,³ despite aggressive campaigns since 2012 to find cases. Let's just go ahead and test our baby boomers.

*Anne C Spaulding, Lesley S Miller

Department of Epidemiology, Rollins School of Public Health, Emory University, Atlanta, GA, USA (ACS); and Department of Medicine, Emory University School of Medicine, Atlanta GA, USA (ACS, LSM)
aspauld@emory.edu

ACS has received grants and personal fees from Gilead Sciences and grants from Bristol-Myers Squibb. LSM has received funding from Bristol-Myers Squibb and grants from Gilead Sciences. They both have received grants from and worked on various projects with members of the Centers for Disease Control and Prevention Division of Viral Hepatitis.

- 1 Joy JB, McCloskey RM, Nguyen T, et al. The spread of hepatitis C virus genotype 1a in North America: a retrospective phylogenetic study. *Lancet Infect Dis* 2016; published online March 30. [http://dx.doi.org/10.1016/S1473-3099\(16\)00124-9](http://dx.doi.org/10.1016/S1473-3099(16)00124-9).

- 2 Armstrong GL, Wasley A, Simard EP, et al. The prevalence of hepatitis C virus infection in the United States, 1999 through 2002. *Ann Intern Med* 2006; **144**: 705–14.
- 3 Centers for Disease Control and Prevention. Hepatitis C: why baby boomers should get tested (factsheet). <http://www.cdc.gov/knowmorehepatitis/Media/PDFs/FactSheet-Boomers.pdf> (accessed March 12, 2016).
- 4 Drummond AJ, Rambaut A, Shapiro B, Pybus OG. Bayesian coalescent inference of past population dynamics from molecular sequences. *Mol Biol Evol* 2005; **22**: 1185–92.
- 5 Larney S, Mahowald MK, Scharff N, Flanigan TP, Beckwith CG, Zaller ND. Epidemiology of hepatitis C virus in Pennsylvania State prisons, 2004–2012: limitations of 1945–1965 birth cohort screening in correctional settings. *Am J Public Health* 2014; **106**: e69–74.
- 6 Moyer VA. Screening for hepatitis C virus infection in adults: U.S. Preventive Services Task Force Recommendation Statement. *Ann Intern Med* 2013; **159**: 349–57.
- 7 Centers for Disease Control and Prevention. Testing recommendations for hepatitis C virus infection. <http://www.cdc.gov/hepatitis/hcv/guidelines.htm> (accessed March 12, 2016).
- 8 Smith BD, Morgan RL, Beckett GA, et al. Recommendations for the identification of chronic hepatitis C virus infection among persons born during 1945–1965. *MMWR Recomm Rep* 2012; **61**: 1–32.
- 9 Shiffman ML. Universal screening for chronic hepatitis C virus. *Liver Int* 2016; **36**: 62–66.
- 10 Cresswell FV, Fisher M, Hughes DJ, Shaw SG, Homer G, Hassan-Ibrahim MO. Hepatitis C core antigen testing: a reliable, quick, and potentially cost-effective alternative to hepatitis C polymerase chain reaction in diagnosing acute hepatitis C virus infection. *Clin Infect Dis* 2015; **60**: 263–66.

Long-term surveillance of the effect of PCV13: the future challenge in Africa



Pneumococcal infections are a leading cause of morbidity and mortality in children. In Africa, the greatest burden of pneumococcal disease and most associated deaths occur in children younger than 5 years.¹ In 2008, this continent still accounted for more than half of the 541 000 deaths caused by pneumococcal disease in this age group.²

The effectiveness of the seven-valent pneumococcal conjugate vaccine (PCV7) has been shown in many countries—the incidence of invasive pneumococcal disease has fallen in all age groups, especially in children younger than 2 years.³ However, in some countries, this decrease was mitigated by an increased incidence of invasive pneumococcal disease caused by non-vaccine serotypes, most belonging to the extended-spectrum PCV.⁴ Soon after the switch from PCV7 to PCV13, many countries such as the USA, France, the UK, and Israel reported the expected pronounced decrease in invasive pneumococcal disease.^{5–8} The high effectiveness of the PCV10 vaccine has also been shown in Finland for invasive

pneumococcal disease and clinically suspected invasive pneumococcal disease.⁹

These results led to PCVs being classified among the more potent available vaccines: in the USA, 55 000 deaths from pneumococcus-related diseases in children were prevented by PCVs between 2001 and 2013.¹⁰ However, in developed countries, which benefit from favourable living conditions and well organised health-care systems, the early (and sustained) assessment of PCV performance is helped by active and well established surveillance of invasive pneumococcal disease. The burden of invasive pneumococcal disease remains highest in low-income countries. Some had introduced PCVs through the GAVI Alliance with a small self-financed contribution. However, little published evidence is available about the effect of PCVs on invasive pneumococcal disease in this context, because of the challenges in surveillance.

In *The Lancet Infectious Diseases*, Grant Mackenzie and colleagues report the results of a well designed population-based active and standardised surveillance study for invasive pneumococcal disease in



Shahzad Noorani/AB/Press Association Images

Published Online
February 17, 2016
[http://dx.doi.org/10.1016/S1473-3099\(16\)00059-1](http://dx.doi.org/10.1016/S1473-3099(16)00059-1)
See [Articles](#) page 703

The Gambia before and after PCV implementation (PCV7 in 2009 and PCV13 in 2011).¹¹ Apart from the reported expected findings, of significance is the admirable effort by the authors to do such a study by rigorously standardising the clinical and laboratory aspects of the surveillance. The project involved several teams, with conceptualisation beginning about 10 years ago. The authors investigated 14 650 patients and identified 320 cases of invasive pneumococcal disease, to do a robust assessment of the effect of PCV13. Despite suboptimal PCV13 coverage, in children 2–23 months of age, the incidence of invasive pneumococcal disease fell by 55% (95% CI 30–71) and that of PCV13 serotype prevalence fell by 82% (95% CI 64–91). In older children (2–4 years of age), the incidence of invasive pneumococcal disease fell by 56% (95% CI 25–75) owing to a 68% (95% CI 39–83) reduction in the prevalence of PCV13 serotypes, which suggests that a combination of direct and herd protection effects were in force.

All these results support the substantial benefit of PCVs introduced in African countries. However, two concerns draw our attention: first, the report of a slight increase in invasive pneumococcal disease incidence in all age groups (except adults) in 2014, and second, a significant increase in prevalence of non-PCV13 serotypes, especially in children aged 2–59 months, in whom the incidence of invasive pneumococcal disease almost doubled. This potentially worrying result should be balanced by the fact that the serotypes that emerged are unlikely to be as invasive as those included in the PCVs.¹² In the post-PCV13 era, the main non-PCV13 serotypes that the authors reported in children 2–59 months old were serotypes 12F, 10A, 15A, 35B, and 16F.¹¹ A recent study showed that of the non-PCV13 serotypes, only two—24F and 12F—had high disease potential in the PCV13 era.¹² In Mackenzie and colleagues' study in The Gambia, serotype 24F was rare (only two cases were reported) in the PCV13 era, so serotype 12F remained the only one with high disease-causing potential.¹¹

Besides the reduced incidence of invasive pneumococcal disease in developed countries, the spectrum and profile of invasive pneumococcal disease has changed following the changed serotype distribution, and the proportion of patients with underlying illnesses has increased.¹³ We have shown that patients

with meningitis are infected more frequently with non-PCV13 serotypes than other serotypes, whereas serotypes implicated in pneumococcal pneumonia are usually PCV13 serotypes, especially 1, 7F, 19A, and 3.¹³ These results agree with those reported by Kaplan and colleagues,¹⁴ showing greater reduction in cases of pneumococcal bacteraemia, pneumonia, and mastoiditis than those of meningitis after PCV13 introduction in eight children's hospitals in the USA. Similarly, the data reported by Mackenzie and colleagues¹¹ show that patients with meningitis were infected more frequently with non-PCV13 serotypes than PCV13 serotypes (44.7% vs 26.3%), whereas for pneumonia, the rates were reversed (21.7% vs 56.1%).¹¹ This serotype distribution clearly explains the notable decrease in pneumonia cases and the slight fall in meningitis cases. With a more prolonged period after PCV13 introduction, the expected decrease in pneumonia cases will be substantial.

This study has important implications for the introduction of PCVs in all African countries, where the rates of invasive pneumococcal disease remain around ten-fold higher than those in developed countries. However, maintenance of invasive pneumococcal disease surveillance is essential to clarify the role of PCVs in changing the serotype incidence, in view of the possible increase in invasive pneumococcal disease cases related to non-vaccine serotypes.

*Corinne Levy, Robert Cohen

Université Paris Est, IMRB- GRC GEMINI, 94000 Créteil, France (CL, RC); ACTIV, Association Clinique et Thérapeutique Infantile du Val de Marne, 27 rue Inkermann, F94100 Saint-Maur des Fossés, France (CL, RC); and Clinical Research Center (CRC), Centre Hospitalier Intercommunal de Créteil, Créteil, France (CL, RC) corinne.levy@activ-france.fr

The authors' institution (ACTIV) has received research grant support from Pfizer, GlaxoSmithKline, and Sanofi Pasteur MSD. CL has received personal fees from Pfizer and Novartis outside the submitted work. RC has received personal fees from Pfizer, GSK, Sanofi, and Novartis outside the submitted work.

Copyright © Levy et al. Open Access article distributed under the terms of CC BY.

- 1 O'Brien KL, Wolfson LJ, Watt JP, et al. Burden of disease caused by *Streptococcus pneumoniae* in children younger than 5 years: global estimates. *Lancet* 2009; **374**: 893–902.
- 2 World Health Organization. Estimated Hib and pneumococcal deaths for children under 5 years of age, 2008 (updated December, 2013). http://www.who.int/immunization/monitoring_surveillance/burden/estimates/Pneumo_hib/en/ (accessed Jan 27, 2016).
- 3 Centers for Disease Control and Prevention. Direct and indirect effects of routine vaccination of children with 7-valent pneumococcal conjugate vaccine on incidence of invasive pneumococcal disease—United States, 1998–2003. *MMWR Morb Mortal Wkly Rep* 2005; **54**: 893–97.

- 4 Miller E, Andrews NJ, Waight PA, Slack MP, George RC. Herd immunity and serotype replacement 4 years after seven-valent pneumococcal conjugate vaccination in England and Wales: an observational cohort study. *Lancet Infect Dis* 2011; **11**: 760–68.
- 5 Ben-Shimol S, Greenberg D, Givon-Lavi N, et al. Early impact of sequential introduction of 7-valent and 13-valent pneumococcal conjugate vaccine on IPD in Israeli children <5 years: an active prospective nationwide surveillance. *Vaccine* 2014; **32**: 3452–59.
- 6 Lepoutre A, Varon E, Georges S, et al. Impact of the pneumococcal conjugate vaccines on invasive pneumococcal disease in France, 2001–2012. *Vaccine* 2015; **33**: 359–66.
- 7 Moore CE, Paul J, Foster D, et al. Reduction of invasive pneumococcal disease 3 years after the introduction of the 13-valent conjugate vaccine in the Oxfordshire region of England. *J Infect Dis* 2014; **210**: 1001–11.
- 8 Simonsen L, Taylor RJ, Schuck-Paim C, Lustig R, Haber M, Klugman KP. Effect of 13-valent pneumococcal conjugate vaccine on admissions to hospital 2 years after its introduction in the USA: a time series analysis. *Lancet Respir Med* 2014; **2**: 387–94.
- 9 Palmu AA, Jokinen J, Nieminen H, et al. Vaccine effectiveness of the pneumococcal *Haemophilus influenzae* protein D conjugate vaccine (PHiD-CV10) against clinically suspected invasive pneumococcal disease: a cluster-randomised trial. *Lancet Respir Med* 2014; **2**: 717–27.
- 10 Whitney CG, Zhou F, Singleton J, Schuchat A, Centers for Disease Control and Prevention. Benefits from immunization during the vaccines for children program era—United States, 1994–2013. *MMWR Morb Mortal Wkly Rep* 2014; **63**: 352–55.
- 11 Mackenzie GA, Hill PC, Jeffries DJ, et al. Effect of the introduction of pneumococcal conjugate vaccination on invasive pneumococcal disease in The Gambia: a population-based surveillance study. *Lancet Infect Dis* 2016; published online Feb 17. [http://dx.doi.org/10.1016/S1473-3099\(16\)00054-2](http://dx.doi.org/10.1016/S1473-3099(16)00054-2).
- 12 Varon E, Cohen R, Bechet S, Doit C, Levy C. Invasive disease potential of pneumococci before and after the 13-valent pneumococcal conjugate vaccine implementation in children. *Vaccine* 2015; **33**: 6178–85.
- 13 Levy C, Varon E, Bechet S, Cohen R. Effect of the 13-valent pneumococcal conjugate vaccine on pneumococcal meningitis in children. *Clin Infect Dis* 2016; **62**: 131–32.
- 14 Kaplan SL, Barson WJ, Lin PL, et al. Early trends for invasive pneumococcal infections in children after the introduction of the 13-valent pneumococcal conjugate vaccine. *Pediatr Infect Dis J* 2013; **32**: 203–07.

The elusive global burden of dengue



Dengue has emerged in the past two decades as a rapidly growing and widespread public health problem, with over half of the world's countries and people now at risk.¹ In new estimates from the Global Burden of Disease study published in *The Lancet Infectious Diseases*, Jeffrey Stanaway and colleagues² suggest that dengue incidence has increased six-fold from 1990 to 2013, accompanied by much flatter mortality trends. Dengue is still regarded as a neglected disease,³ yet its incidence is increasing at an alarming rate, by contrast with declines in other neglected diseases.⁴ WHO estimates that 50 million to 100 million cases occur annually.⁵ In

2012, however, formal modelling put the number of dengue infections as high as 390 million (95% credible interval 284 million–528 million), but included both inapparent and apparent cases.¹ Their estimated number of apparent cases (96 million, 95% credible interval 67 million–136 million) lies at the high end of the WHO estimates.

Against this background, these new estimates from the Global Burden of Disease Study suggest a much lower number of cases for 2013: 58.4 million (95% uncertainty interval 23.6 million–121.9 million).² Why is it so difficult to arrive at consistent estimates?

Published Online
February 10, 2016
[http://dx.doi.org/10.1016/S1473-3099\(16\)00076-1](http://dx.doi.org/10.1016/S1473-3099(16)00076-1)
See [Articles](#) page 712




Figure: Dengue case fatality rates for 39 countries estimated to have at least ten dengue deaths and autochthonous transmission
Data from Stanaway and colleagues.² CFR=case fatality rate.